

Science & Healthcare


PEER-REVIEWED MEDICAL SCIENTIFIC JOURNAL

**РЕЦЕНЗИРУЕМЫЙ МЕДИЦИНСКИЙ
НАУЧНО-ПРАКТИЧЕСКИЙ ЖУРНАЛ**


Наука и Здравоохранение

Ғылым және Денсаулық Сақтау

**РЕЦЕНЗИЯЛАНАТЫН МЕДИЦИНАЛЫҚ
ҒЫЛЫМИ-ПРАКТИКАЛЫҚ ЖУРНАЛ**



**Volume 26,
Issue 5
2024**



Министерство здравоохранения
Республики Казахстан
Учредитель: НАО
«Медицинский университет Семей»
Основан в 1999 году.

Журнал зарегистрирован в
Министерстве информации и
коммуникаций Республики Казахстан
Комитете информации № 17773-Ж.

Входит в Перечень изданий,
рекомендуемых Комитетом по
обеспечению качества в сфере науки и
высшего образования МНВО
Республики Казахстан для публикации
основных результатов научной
деятельности (Приказ №603 от
12.07.2024г.)

Включен в Ulrich's Periodicals
Directory, Global Health, CAB
Abstracts, InfoBase Index, Directory
of Research Journals Indexing,
Российский индекс научного
цитирования (РИНЦ), E-library.ru,
Cyberleninka.ru, Norwegian register
for scientific journals (NSD),
Всесоюзный институт научной и
технической информации Российс-
кой академии наук (ВИНИТИ РАН),
Ассоциацию CONEM, РАЦС,
DataBase Indexing, ICI World of
Journals

Подписной индекс 74611
в каталоге «Казпочта»
Цена свободная

Сайт <http://newjournal.ssmu.kz>
e-mail: selnura@mail.ru

Адрес редакции:

071400, г. Семей
ул. Абая Кунанбаева, 103
контактный телефон:
(7222) 56-42-09 (вн. № 1054)
факс: (7222) 56-97-55

Выпускающий редактор:

Э.Ф. Сапаргалиева

Переводчики:

С.А. Жаукенова, Н.А. Шумский



При цитировании материалов
ссылка на журнал обязательна.

Отпечатано в типографии
Медицинского университета Семей
Подписано в печать: 30.10.2024г.
Формат 60х90/8.

Печать цифровая. Усл.п.л 30,8
Тираж 500 экз., зак.184

Online ISSN: 3007 – 0244

Print ISSN: 2410 – 4280

НАУКА И ЗДРАВООХРАНЕНИЕ

РЕЦЕНЗИРУЕМЫЙ МЕДИЦИНСКИЙ
НАУЧНО-ПРАКТИЧЕСКИЙ ЖУРНАЛ

Том 26 №5, 2024

Журнал «Наука и Здравоохранение» - рецензируемый медицинский научно-практический журнал, который публикует результаты оригинальных исследований, обзоры литературы, клинические случаи, краткие сообщения и отчеты о конференциях по широкому кругу вопросов, связанных с клинической медициной и общественным здоровьем. Основной читательской аудиторией журнала является медицинское научное сообщество, практикующие врачи, докторанты и магистранты в области медицины и общественного здоровья.

Главный редактор:

А.А. Дюсупов

доктор медицинских наук, профессор

Зам. главного редактора:

Д.М. Шабдарбаева

доктор медицинских наук, профессор

Редакционный совет:

Абдрахманов А.С. (Нур-Султан, Казахстан)
Акильжанова А.Р. (Нур-Султан, Казахстан)
Акшулаков С.К. (Нур-Султан, Казахстан)
Баймаханов Б.Б. (Алматы, Казахстан)
Брузати Лука Джиованни Карло (Удин, Италия)
Виджай Кумар Чатту (Торонто, Канада)
Гржибовский А.М. (Архангельск, Российская Федерация)
Гюрель Фазыл Сердар (Анкара, Турция)
Даутов Т.Б. (Нур-Султан, Казахстан)
Джерзи Крупински Белецки (Барселона, Испания)
Жумадилов Ж.Ш. (Нур-Султан, Казахстан)
Кавальчи Чемиль (Анкара, Турция)
Карпенко А.А. (Новосибирск, Российская Федерация)
Ковальчук В.В. (Санкт-Петербург, Российская Федерация)
Лесовой В.Н. (Харьков, Украина)
Лукьянов С.А. (Москва, Российская Федерация)
Мутиг К. (Шарите, Германия)
Носо Й. (Шимане, Япония)
Раманкулов Е.М. (Нур-Султан, Казахстан)
Степаненко В.Ф. (Обнинск, Российская Федерация)
Тапбергенов С.О. (Семей, Казахстан)
Тринчеро Элизабетта Флора Ольга (Милан, Италия)
Хоссейни Хенгаме (Скрантон, Соединенные Штаты Америки)
Хоши М. (Хиросима, Япония)
Шейнин А. (Тель-Авив, Израиль)

The Ministry of Healthcare
of the Republic of Kazakhstan
Publisher: NCJSC
«Semey Medical University»
Established in 1999

Journal is registered in Ministry of Information and Communication of the Republic of Kazakhstan in the Committee of Information. Certificate of registration of a periodical printed publication № 17773-Ж.

The journal is included in the List of publications recommended by the Committee for Quality Assurance in the Field of Science and Higher Education of the Ministry of Science and Higher Education of the Republic of Kazakhstan for the publication of the main results of scientific activity (Order from 12.07.2024 №603)

The journal is indexed in Ulrich's Periodicals Directory, Global Health, CAB Abstracts, InfoBase Index, Directory of Research Journals Indexing, Russian Science Citation Index, Scientific electronic library E-library.ru, Cyberleninka.ru, NSD (Norwegian register for scientific journals), VINITI RAN, CONEM, RASS, DataBase Indexing, ICI World of Journals

Subscription index in catalogue of "Kazpost" 74611

Open price.

Website <http://newjournal.ssmu.kz>

e-mail: selnura@mail.ru

Address of editor office and publisher:

071400, Semey, Abay st. 103,
Tel. (7222) 56-42-09 (in1054)
Fax: (7222) 56-97-55

Publishing editor:

E.F. Sapargaliyeva

Translators:

S.A. Zhaukenova,
N.A. Shumskiy



In case of citation of materials, a link on the journal is required.

Printed by printing office of Semey medical university

Signed in press on 28 October 2024

Format 60x90/8. 30,8 Digital printing.

Circulation 500 copies, ord. 184

Online ISSN: 3007 – 0244

Print ISSN: 2410 – 4280

SCIENCE & HEALTHCARE

PEER-REVIEWED SCIENTIFIC-PRACTICAL
MEDICAL JOURNAL

Volume 26 №5, 2024

«Science & Healthcare» is a peer-reviewed scientific-practical medical journal, which publishes original articles, literature reviews, clinical case, short communications and conference reports covering all areas of clinical medicine and public health. The primary audience of the journal includes medical scientific community, practicing physicians, doctoral- and master - students in the fields of medicine and public health.

Chief editor:

A.A. Dyussupov

Doctor of medical science, Professor

Deputy Editor in Chief:

D.M. Shabdarbayeva

Doctor of medical science, Professor

Editorial board:

Abdrakhmanov A.S. (Nur-Sultan, Kazakhstan)

Akilzhanova A.R. (Nur-Sultan, Kazakhstan)

Akshulakov S.K. (Nur-Sultan, Kazakhstan)

Baimakhanov B.B. (Almaty, Kazakhstan)

Brusati Luca G. (Udine, Italy)

Vijai Kumar Chattu (Toronto, Canada)

Grijbovski A.M. (Arkhangelsk, Russian Federation)

Gurel Fazil Serdar (Ankara, Turkey)

Jerzy Krupinski Bielecki (Barselona, Spain)

Dautov T.B. (Nur-Sultan, Kazakhstan)

Zhumadilov Zh.Sh. (Nur-Sultan, Kazakhstan)

Karpenko A.A. (Novosibirsk, Russian Federation)

Kavalci Cemil (Ankara, Turkey)

Kovalchuk V.V. (St. Petersburg, Russian Federation)

Lesovoy V.N. (Kharkiv, Ukraine)

Luk'yanov S.A. (Moscow, Russian Federation)

Mutig K. (Sharite, Germany)

Noso Y. (Shimane, Japan)

Ramankulov Ye.M. (Nur-Sultan, Kazakhstan)

Stepanenko V.F. (Obninsk, Russian Federation)

Tapbergenov S.O. (Semey, Kazakhstan)

Trincherio Elisabetta Flora Olga (Millan, Italy)

Hosseini Hengameh (Scranton, United States of America)

Hoshi M. (Hiroshima, Japan)

Sheinin Anton (Tel-Aviv, Israel)

Қазақстан Республикасы
денсаулық сақтау министрлігі
Құрылтайшы: КеАҚ
«Семей медицина университеті»
1999 негізі салынды

Журнал Қазақстан Республикасы
ақпарат және коммуникациялар
министрлігі, Ақпарат комитетінде
тіркелген. Мерзімді баспасөз
басылымын есепке қою туралы куәлігі
№ 17773-Ж

Журнал Ғылыми еңбектің
негізгі нәтижелерін жариялау үшін
Қазақстан Республикасы Ғылым
және жоғары білім министрлігінің
Ғылым және жоғары білім
саласында сапаны қамтамасыз ету
комитеті ұсынатын ғылыми
басылымдар Тізбесіне кіреді
(Бұйрық №603 12.07.2024ж.)

Ulrich's Periodicals Directory,
Global Health, CAB Abstracts,
InfoBase Index, Directory of
Research Journals Indexing,
Ғылыми дәйектеу Ресейлік
индекс (РИНЦ), E-library.ru. -
Ғылыми электронды кітапханаға,
Cyberleninka.ru, NSD (Norwegian
register for scientific journals),
ВИНИТИ РАН, CONEM, РАЦС,
DataBase Indexing, ICI World of
Journals, енгізілді.

Каталогтағы жазылу индексі
«Казпочта» 74611

Бағасы еркін

Сайт <http://newjournal.ssmu.kz>

e-mail: selhura@mail.ru

Баспаның және баспагердің мекен-
жайы:

071400, Семей қаласы,

Абай көшесі, 103.

тел. (87222) 56-42-09 (ішкі 1054);

факс: (7222) 56-97-55

Баспа редакторы:

Э.Ф. Сапарғалиева

Аудармашылар:

С.А. Жаукенова, Н.А. Шумский



Материалдарды дәйектеу кезінде
журналға сілтеме жасау міндетті.

Семей медицина университетінің
баспаханасында басылған

Баспаға қол қойылды 30.10.2024.

Формат 60x90/8. Баспа сандық.

Шартты-баспа парағы 30,8

Таралуы 500 дана. Зак.184

Online ISSN: 3007 – 0244

Print ISSN: 2410 – 4280

ҒЫЛЫМ ЖӘНЕ ДЕНСАУЛЫҚ САҚТАУ РЕЦЕНЗИЯЛАНАТЫН МЕДИЦИНАЛЫҚ ҒЫЛЫМИ-ПРАКТИКАЛЫҚ ЖУРНАЛ

Том 26 №5, 2024

«Ғылым және денсаулық сақтау» журналы -
рецензияланатын медициналық ғылыми-практикалық
журнал, клиникалық медицина мен қоғамдық денсаулықпен
байланысты бірегей зерттеулер нәтижелерін, әдеби
шолуларды, кең шеңберлі сұрақтар бойынша
конференциялар туралы қысқа мәлімдемелер мен есептерді
жариялайды. Медициналық ғылыми қоғамдастық,
тәжірибелік дәрігерлер, медицина мен қоғамдық денсаулық
саласындағы докторанттар мен магистранттар журналдың
негізгі оқырман аудиториясы болып табылады.

Бас редактор:

А.А. Дүсіпов

медицина ғылымдарының докторы, профессор

Бас редактордың орынбасары: Д.М. Шабдарбаева,

медицина ғылымдарының докторы, профессор

Редакциялық кеңес:

Абдрахманов А.С. (Нұр-Сұлтан, Қазақстан)

Ақылжанова А.Р. (Нұр-Сұлтан, Қазақстан)

Акшулаков С.К. (Нұр-Сұлтан, Қазақстан)

Баймаханов Б.Б. (Алматы, Қазақстан)

Брузати Лука Джиованни Карло (Удин, Италия)

Виджай Кумар Чатту (Торонто, Канада)

Гржибовский А.М. (Архангельск, Российская Федерация)

Гюрель Фазыл Сердар (Анкара, Түркия)

Джерзи Крупински Белецки (Барселона, Испания)

Даутов Т.Б. (Нұр-Сұлтан, Қазақстан)

Жумадилов Ж.Ш. (Нұр-Сұлтан, Қазақстан)

Кавальчи Чемиль (Анкара, Түркия)

Карпенко А.А. (Новосибирск, Ресей Федерациясы)

Ковальчук В.В. (Санкт-Петербург, Ресей Федерациясы)

Лесовой В.Н. (Харьков, Украина)

Лукиянов С.А. (Москва, Ресей Федерациясы)

Мутиг К. (Шарите, Германия)

Носо Й. (Шимане, Жапония)

Раманқұлов Е.М. (Нұр-Сұлтан, Қазақстан)

Степаненко В.Ф. (Обнинск, Ресей Федерациясы)

Тапбергенов С.О. (Семей, Қазақстан)

Тринчеро Элизабетта Флора Ольга (Милан, Италия)

Хоссейни Хенгаме (Скрантон, Америка Құрама Штаттары)

Хоши М. (Хиросима, Жапония)

Шейнин А. (Тель-Авив, Израиль)

СОДЕРЖАНИЕ

- Оразбаева Д.Р., Иса Г., Сагындыков И.К., Ахметов Е.А., Манатова А.М., Жарлыганова Д.С.**
Роль ультразвукового исследования в выявлении сосудистых осложнений у реципиентов трансплантата печени
- Skenderova A.A., Nuskabayeva G.N., Tatykayeva U.B., Sarsenbayeva G.Zh., Kemelbekov K.S., Seidakhmetova A.A.**
Therapeutic effects of melatonin in an ethanol-induced gastric ulcer model
- Bedel H.-A., Bedel C., Selvi F., Zortuk Ö., Karancı Yu.**
Effect of Seasonal Changes on scorpion poison
- Абуова А.С., Мейрамова Ә.М., Абдахина Б.Б., Бокаева А.А., Цой Т.Т., Бикташев Д.Б., Айнабекова Б.А.**
Особенности течения анкилозирующего спондилоартрита в казахстанской популяции в зависимости от гендерной принадлежности
- Apbassova M., Apbassova S.A., Shabdarbayeva D.M., Chaizhunusova N.Zh., Bulegenov T.A., Mutig K., Mussina A.A., Baurzhan A.B., Assubayev O.B., Dyusembayev A.K., Lepesbayev M.N., Amantayeva G.K., Tatykayeva U.B., Svitich T.N., Shalgumbayeva G.M., Dyussupov A.A.**
Suicide rate forecasting in the cities of Ust-Kamenogorsk, Semey, and Pavlodar region using a mathematical model
- Byltash S., Brimzhanova M., Faizullina K., Akhtayeva N., Alekenova N., Kosherbayeva L., Nurbaulina E.**
Current practice in caring for people with dementia
- Akylzhanova Zh.E., Tanysheva G.A., Gasanova E.Z., Bilyalova G.T., Nurmagambetova D.M., Akhmetova A.B., Suleimenova A.P., Ulan S.M., Nurmakhan N.S.**
Using the Robson classification to estimate the caesarean section rate and look for ways to reduce it

COVID-19 - АКТУАЛЬНАЯ ТЕМА

- Дюсупов А.А., Жуманбаева Ж.М., Кайрханова Ы.О., Досбаева А.Д., Крыкпаева А.С., Серикбаев А.С., Амренова К.Ш., Сеиткабылов А.А., Кудайбергенов Р.Н.**
Пост-COVID-19-синдром: симптомы и факторы риска

ОБЗОР ЛИТЕРАТУРЫ

- Shokenov R.D., Shakhmarova T.K., Mirmanova Zh.Zh., Chamoieva A.Ye., Zhalbinova M.R., Rakhimova S.E., Bekbossynova M.S., Akilzhanova A.R.**
Characterizing the Role of ABCG5/G8 in Sitosterolemia: Diagnostic Challenges in Differentiating from Familial Hypercholesterolemia
- Samatova N.E., Kamalbekova G.M., Jaxalykova K.K., Mussina A.A., Karin B.T.**
The main tools of the confidential audit of perinatal mortality: a review

- 79-84 **Orazbaeva D.R., Isa G., Sagyndykov I.K., Akhmetov E.A., Manatova A.M., Zharlyganova D.S.**
The role of ultrasound in identifying vascular complications in liver transplant recipients
- 85-92 **Скендерова А.А., Нускабаева Г.О., Татыкаева У.Б., Сарсенбаева Г.Ж., Кемельбеков К.С., Сейдахметова А.А.**
Терапевтические эффекты мелатонина на модели язвы желудка, вызванной этанолом
- 93-98 **Бедел Х.-А., Бедел Дж., Сельви Ф., Зортук О., Каранджи Ю.**
Влияние сезонных изменений на яд скорпиона
- 99-107 **Abuova A.S., Meiramova A.M., Abdakhina B.B., Bokayeva A.A., Tsoy T.T., Biktashev D.B., Ainabekova B.A.**
Features of the ankylosing spondylitis course depending on gender in the Kazakh population
- 108-114 **Апбасова М., Апбасова С.А., Шабдарбаева Д.М., Чайжунусова Н.Ж., Булегенов Т.А., Мутиг К., Мусина А.А., Бауржан А.Б., Асубаев О.Б., Дюсембаев А.К., Лепесбаев М.Н., Амантаева Г.К., Татыкаева У.Б., Свитич Т.Н., Шалгумбаева Г.М., Дюсупов А.А.**
Прогнозирование уровня суицида в городах Усть-Каменогорск, Семей и Павлодарской области с помощью математической модели
- 115-120 **Былташ С., Бримжанова М., Файзуллина К., Ахтаева Н., Алеkenова Н., Кошербаева Л., Нурбаулина Э.**
Существующая практика ухода за людьми с деменцией
- 121-128 **Акылжанова Ж.Е., Танышева Г.А., Гасанова Э.З., Билялова Г.Т., Нурмагамбетова Д.М., Ахметова А.Б., Сулейменова Ә.С., Ылан С.М., Нурмахан Н.С.**
Применение классификации Робсона для оценки показателя частоты кесарева сечения и поиска возможностей его снижения

COVID-19 - TOPICAL SUBJECT

- 129-140 **Dyussupov A.A., Zhumanbayeva Zh.M., Kairkhanova Y.O., Dosbayeva A.M., Krykpaeva A.S., Serikbayev A.S., Amrenova K.Sh., Seitkabylov A.A., Kudaibergenov R.N.**
Post-COVID-19 syndrome: symptoms and risk factors

REVIEWS

- 141-149 **Шокенов Р.Д., Шахмарова Т.К., Мирманова Ж.Ж., Чамойева А.Е., Жалбинова М.Р., Рахимова С.Е., Бекбосынова М.С., Акильжанова А.Р.**
Характеристика роли ABCG5/G8 при ситостеролемии: диагностические трудности в дифференциации от семейной гиперхолестеринемии
- 150-159 **Саматова Н.Е., Камалбекова Г.М., Джаксалыкова К.К., Мусина А.А., Карин Б.Т.**
Основные инструменты конфиденциального аудита перинатальной смертности: обзор литературы

Received: 08 June 2024 / Accepted: 14 October 2024 / Published online: 31 October 2024

DOI 10.34689/SH.2024.26.5.011

UDC 616.33-002.44-085 : 612.43



THERAPEUTIC EFFECTS OF MELATONIN IN AN ETHANOL-INDUCED GASTRIC ULCER MODEL

Akbota A. Skenderova¹, <https://orcid.org/0000-0002-5392-9433>

Gulnaz O. Nuskabayeva¹, <https://orcid.org/0000-0002-2906-1427>

Ugilzhan B. Tatykayeva¹, <https://orcid.org/0000-0002-7519-479X>

Gulzat Zh. Sarsenbayeva², <https://orcid.org/0000-0002-1518-6528>

Kanatzhan S. Kemelbekov², <https://orcid.org/0009-0008-1619-6278>

Aizat A. Seidakhmetova², <https://orcid.org/0000-0002-5794-3814>

¹ Akhmet Yassawi International Kazakh-Turkish University, Turkistan, Republic of Kazakhstan;

² South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan.

Abstract

Introduction. Recently, melatonin and its agonists have been proven to be effective in the treatment of jet lag syndrome, insomnia, stress, fear, anger, anxiety, stress conditions, which are accompanied by sleep disorders, increase the immune system of the whole body, and as antioxidant protection. Melatonin is known to be an important chronobiological molecule that exerts its effect on any system in the body. The role and pharmacological effect of melatonin in the gastrointestinal system is still one of the important topics, and research continues in various directions.

Aim. To monitor the therapeutic effects of melatonin in relation to ethanol concentration in an ethanol-induced experimental gastric ulcer model.

Material's and method. A prospective preclinical (experimental) single-center study. The work was carried out using laboratory rats. The research work was carried out for each of the 6 groups, in a model of gastric ulcer induced by ethanol of various concentrations (100-75-60%) and melatonin was administered to the ulcer (15 mg/kg) and the control group. The obtained data were subjected to statistical analysis using a trial version of the SPSS statistical program (SPSS 20 Inc, Chicago, IL, USA), the critical significance level (p) was assumed to be 0.05.

Results. In addition to this increase in the wound area, melatonin administered as a prophylactic subcutaneous injection reduced the overall wound area. In the group of wounds induced by ethanol (100%), the wound area was 15.01 ± 1.92 mm, after intraperitoneal injection of melatonin (15 mg/kg), and after 30 minutes the wound area was 14.01 ± 1.74 mm in the 100% ethanol-induced group and 13.24 ± 1.71 mm, in the 75% ethanol-induced group, 11.87 ± 1.45 mm in the melatonin + 75% ethanol-induced group, 11.52 ± 1.15 mm in the 60% ethanol-induced group and 6.12 ± 0.54 mm in the melatonin + 60% ethanol-induced group.

Conclusion. In this study, melatonin showed a positive effect in the prevention and treatment of ethanol-induced ulcers at different concentrations. A significant change was observed at higher levels of melatonin inhibition of ethanol exposure at lower concentrations.

Keywords: melatonin, antioxidant, circadian rhythm, metabolic syndrome, digestive system, gastrointestinal ulcer.

Резюме

ТЕРАПЕВТИЧЕСКИЕ ЭФФЕКТЫ МЕЛАТОНИНА НА МОДЕЛИ ЯЗВЫ ЖЕЛУДКА, ВЫЗВАННОЙ ЭТАНОЛОМ

Ақбота А. Скендерова¹, <https://orcid.org/0000-0002-5392-9433>

Гульназ О. Нускабаева¹, <https://orcid.org/0000-0002-2906-1427>

Угилжан Б. Татыкаева¹, <https://orcid.org/0000-0002-7519-479X>

Гульзат Ж. Сарсенбаева², <https://orcid.org/0000-0002-1518-6528>

Канатжан С.Кемельбеков², <https://orcid.org/0009-0008-1619-6278>

Айзат А. Сейдахметова², <https://orcid.org/0000-0002-5794-3814>

¹ Международного казахско-турецкого университета им. Х.А. Ясави, г. Туркестан, Республика Казахстан;

² АО «Южно-Казахстанская медицинская академия», г. Шымкент, Республика Казахстан.

Актуальность. В последнее время мелатонин и его агонисты доказали свою эффективность в качестве антиоксидантной защиты при лечении синдрома смены биологических ритмов, бессонницы, стресса, страха, гнева, тревоги, стрессовых состояний, которые сопровождаются нарушениями сна и повышают иммунитет всего организма. Известно, что мелатонин является важной хронобиологической молекулой, оказывающей влияние на любую систему организма. Роль и фармакологический эффект мелатонина в системе желудочно-кишечного тракта до сих пор остается одной из важных тем и исследования продолжают в различных направлениях.

Цель. Исследование терапевтических эффектов мелатонина в зависимости от концентрации этанола в модели

Материалы и методы. Проспективное доклиническое (экспериментальное) одноцентровое исследование. Работа проводилась с использованием лабораторных крыс. Исследовательская работа проводилась для каждой из 6 групп, в модели язвы желудка, индуцированной этанолом различной концентраций (100-75-60%) и вводился мелатонин до язвы (15 мг/кг) и контрольной группе. Полученные данные были подвергнуты статистическому анализу с помощью пробной версии статистической программы SPSS (SPSS 20 Inc, Chicago, IL, USA), критический уровень значимости (p) принимался за 0,05.

Результаты. В желудке крыс этанол при макроскопическом анализе в зависимости от концентрации увеличивал площадь раны. Мелатонин вводимый в виде профилактической подкожной инъекции, уменьшал общую площадь раны. В группе индуцированным этанолом (100%), площадь раны без введения мелатонина составила $15,01 \pm 1,92$ мм., а с введением мелатонина (15 мг/кг) до язвы площадь раны уменьшилась на $14,01 \pm 1,74$ мм., и в группе 75% этанола, площадь раны без введения мелатонина составила $13,24 \pm 1,71$ мм., введением мелатонина до язвы площадь раны уменьшилась на $11,87 \pm 1,45$ мм., и в группе 60% этанола, площадь раны без введения мелатонина составила $11,52 \pm 1,15$ мм., и введением мелатонина до язвы площадь раны уменьшилась на $6,12 \pm 0,54$ мм.

Выводы. В этом исследовании мелатонин показал положительный эффект в профилактике и лечении язв, вызванных этанолом, в различных концентрациях. Значительное изменение наблюдалось в более высоких уровнях ингибирования мелатонина при воздействии этанола в более низких концентрациях.

Ключевые слова: мелатонин, антиоксидант, циркадный ритм, метаболический синдром, пищеварительная система, язва желудочно-кишечного тракта.

Түйіндеме

ЭТАНОЛМЕН ШАҚЫРЫЛҒАН АСҚАЗАН ЖАРА МОДЕЛІНДЕ МЕЛАТОНИННІҢ ЕМДІК ӘСЕРІ

Ақбота А. Скендерова¹, <https://orcid.org/0000-0002-5392-9433>

Гульназ О. Нускабаева¹, <https://orcid.org/0000-0002-2906-1427>

Угилжан Б. Татыкаева¹, <https://orcid.org/0000-0002-7519-479X>

Гульзат Ж. Сарсенбаева², <https://orcid.org/0000-0002-1518-6528>

Канатжан С.Кемельбеков², <https://orcid.org/0009-0008-1619-6278>

Айзат А. Сейдахметова², <https://orcid.org/0000-0002-5794-3814>

¹ Қ.А. Ясауи атындағы Халықаралық қазақ-түрік университеті, Түркістан қ., Қазақстан Республикасы;

² «Оңтүстік Қазақстан медицина академиясы» АҚ, Шымкент қ., Қазақстан Республикасы.

Өзектілігі. Қазіргі таңда, мелатонин және агонистерін тек ұйқының бұзылуымен бірге жүретін жетлаг синдромы, ұйқысыздық, стресс, қорқыныш, ашулану, мазасыздық, күйзелістік жағдайларды емдеуде, жалпы ағзаның иммундық жүйесін көтеруде, антиоксиданттық қорғаныс ретінде тиімділігі дәлелденген. Мелатонин организмде кез келген жүйеде әсерін көрсететін маңызды хронобиологиялық молекула екендігі бұрыннан мәлім. Мелатониннің асқазан-ішек жүйесіндегі рөлі және фармакологиялық әсері осы күнге дейін өзекті тақырыптардың бірі болып, түрлі бағыттарда зерттеулер жалғасын табуда.

Зерттеудің мақсаты. Этанолмен шақырылған экспериментальді асқазан жара моделінде этанол концентрациясына байланысты мелатониннің емдік әсерін бақылау.

Зерттеудің материалдары мен әдістері. Проспективті клиникаға дейінгі (эксперименттік) бір орталықты зерттеу. Жұмыс зертханалық егеуқұйрықтарды қолдану арқылы жүргізілді. Зерттеу жұмыстары 6 топтың әрқайсысы үшін әр түрлі концентрациядағы этанолмен индукцияланған асқазан жарасының моделінде (100-75-60%) жүргізілді және мелатонин ойық жараға (15 мг/кг) және бақылау тобына енгізілді. Алынған мәліметтер SPSS статистикалық бағдарламасының сынақ нұсқасында (SPSS 20 Inc, Chicago, IL, USA) талдаудан өтті, статистикалық маңызды деңгейі $p < 0,05$ деп қабылданды.

Зерттеу нәтижелері. Егеуқұйрық асқазанында этанол концентрация тәуелді макроскопиялық талдауда пайда болған жара ауданын арттырды. Осы жара ауданының артуында мелатонин алдын ала профилактикалық мақсатта теріастылық инъекция түрінде берілгенде, жараның жалпы ауданын азайтқан. Этанолмен (100%) шақырылған жара тобында пайда болған жара ауданы $15,01 \pm 1,92$ мм болса, алдын ала мелатонин (15 мг/кг) перитон ішіне беріліп, 30 минуттан соң 100% этанолмен шақырылған жара тобында $14,01 \pm 1,74$ мм, 75% этанол тобында $13,24 \pm 1,71$ мм болса, мелатонин+ 75% этанолмен шақырылған жара тобында $11,87 \pm 1,45$ мм, 60% этанолмен шақырылған жара тобында $11,52 \pm 1,15$ мм болса, ал мелатонин+ 60% этанолмен шақырылған жара тобында $6,12 \pm 0,54$ мм жара ауданы азайғандығы байқалған.

Қорытындылары. Осы зерттеуде әр түрлі концентрацияда этанолдан пайда болған жараның алдын алуда, емдеуде мелатонин оң әсерін көрсеткен. Неғұрлым төмен концентрациядағы этанолдың әсеріне мелатонин жараның алдын алуда соғұрлым жоғары деңгейде мәнді өзгеріс байқалған.

Түйінді сөздер: Мелатонин, антиоксидант, циркадты ырғақ, метаболикалық синдром асқорыту жүйесі, асқазан-ішек жолдарының жара ауруы.

For citation / Для цитирования / Дәйексөз үшін:

Skenderova A.A., Nuskabayeva G.N., Tatykayeva U.B., Sarsenbayeva G.Zh., Kemelbekov K.S., Seidakhmetova A.A. Therapeutic effects of melatonin in an ethanol-induced gastric ulcer model // *Nauka i Zdravookhranenie* [Science & Healthcare]. 2024. Vol.26 (5), pp. 85-92. doi 10.34689/SH.2024.26.5.011

Скендерова А.А., Нускабаева Г.О., Татыкаева У.Б., Сарсенбаева Г.Ж., Кемельбеков К.С., Сейдахметова А.А. Терапевтические эффекты мелатонина на модели язвы желудка, вызванной этанолом // *Наука и Здоровоохранение*. 2024. Т.26 (5). С. 85-92. doi 10.34689/SH.2024.26.5.011

Скендерова А.А., Нускабаева Г.О., Татыкаева У.Б., Сарсенбаева Г.Ж., Кемельбеков К.С., Сейдахметова А.А. Этанолмен шақырылған асқазан жара моделінде мелатониннің емдік әсері // *Ғылым және Денсаулық сақтау*. 2024. Т.26 (5). Б. 85-92. doi 10.34689/SH.2024.26.5.011

Introduction

Melatonin is a natural neurotransmitter involved in the regulation of many biological and physiological processes, sleep, and biorhythms (circadian rhythms) in the body. Its main task is to control the body's biological clock and to maintain its rhythm [8]. Melatonin (chemical name 5-methoxy N-acetyl tryptamine) is an indole hormone secreted mainly by the pineal gland and synthesized by other peripheral systems such as the brain, bone marrow, platelets, lymphocyte cells, and skin, and affects many systems [12-16]. Its effects on cell regeneration and the immune system, as well as other known functions, are known. This literature review discusses the role of melatonin synthesized in the gastrointestinal (GI) tract for the body and its effects on physiological and pathological processes, focusing on the general characteristics of the hormone melatonin. Melatonin is synthesized from enter chromaffin cells in the gastrointestinal tract, with the intensity of this synthesis coordinated with the rhythm of food intake. The concentration of melatonin in the gastrointestinal tract is about 10-100 times that in blood. In fact, the total amount of melatonin in the gastrointestinal mucosa is 400 times greater than in the pineal gland [18-19].

Synthesis of the hormone melatonin. Tryptophan, a precursor for the synthesis of melatonin, an essential amino acid, is obtained from food. After the removal of tryptophan from the plasma through the pineal gland, 5-hydroxytryptophan, the first intermediate for the synthesis of serotonin and melatonin, is formed in pinealocytes by hydroxylation with tryptophan hydroxylase and easily passes the blood-brain barrier. Using the enzyme tryptophan hydroxylase, tetrahydrobiopterin (BH4) and superoxide (O₂), and vitamin B6 as coenzymes, 5-hydroxytryptophan is converted into 5-hydroxytryptamine (serotonin) by the enzyme 5-hydroxytryptophan decarboxylase. Serotonin cannot cross the blood-brain barrier. Serotonin is then acetylated with N-acetylserotonin by the enzyme N-acetyltransferase (NAT). N-acetyl serotonin combines with HYOMT (hydroxyl indole methyltransferase) to N-acetyl 5-methoxy tryptamine, i.e. it is converted to melatonin [3-5].

Sites of synthesis of the hormone melatonin. In mammals, melatonin is secreted by the pineal gland, the biliary and gastrointestinal tracts, the pigment epithelium, and retinal photoreceptors and plays a role in regulating the retinal response to the change of day and night. Melatonin, produced in the skin, is responsible for tissue changes and protection against harmful sunlight.

After a meal, melatonin, which is synthesized by the chromaffin cells of the gastrointestinal tract, is released into the bloodstream. Melatonin is synthesized in the bile-oxidized cholesterol derivatives and protects the mucous layer and epithelium of the bile ducts from oxidative damage by bile acids. However, this synthesized amount has very little effect on circulating melatonin levels. The amount of melatonin in the circulating pineal gland synthesizes is about 80% [1,6,9].

Melatonin receptors. Three types of melatonin receptors have been identified. Activation of the MT1 receptor exerts its effect by reducing adenylate cyclase activity in cells. This receptor is responsible for the retina, kidney function, biological rhythm, fertility function and contraction of cerebral arteries [2]. On the other hand, MT2 and MT3 receptors act by hydrolyzing phosphoinositol. Melatonin receptors are found in various parts of the human brain, intestine, ovaries and blood vessels. Known agonists are used for sleep disorders and depression: the melatonin analogue ramelteon, tasimelteon, agomelatine [11-12].

Pharmacokinetics. Melatonin binds 60-70% to albumin in the blood. The elimination half-life of melatonin is 3-45 minutes. Melatonin, which is partially contained in water and lipids in large quantities, is easily absorbed into tissues and cells. Melatonin is first converted in the liver to 6-hydroxy dopamine and then to N-acetyl-5-methoxy-6-hydroxytryptamine via a series of reactions. It is then converted to 6-sulphatoxymelatonin, conjugated to sulfate or glucuronide, and excreted in the urine. 1% is excreted unchanged in the urine. Because of age, melatonin levels are low in newborns during the first three months of life. 65 pg/ml at 5 to 15 years of age, 20 pg/ml at 50 to 70 years of age. In adults, average plasma levels are 50-70 pg/ml [23].

Melatonin hormone secretion is dependent on the photosensitivity of pinealocyte cells, light causes synthesis to slow down, and in darkness, and melanocytes begin to secrete melatonin again. Melatonin secretion is highest between 23:00 and 05:00, and the concentration in the blood increases by a factor of 3-10. The release of melatonin has a specific circadian rhythm. It starts increasing in the evening at 21:00-22:00 and peaks at 02:00-04:00. It begins to decrease at 05:00-07:00 in the morning. It starts after 07.00 and decreases to basal levels. The blood concentration of melatonin is 0-20 pg/dl during the day and rises to 50-200 pg/dl at night. An average of 30 mg of melatonin is synthesized during the night [7,10].

The antioxidant effect of melatonin. Oxidative stress caused by certain toxins, which cause oxidative damage to

tissue, is stopped by melatonin. As melatonin is water- and lipid-soluble, it easily penetrates into all intracellular components and effectively protects the cell membrane, organelles, and nucleus from free radical damage. Melatonin on the outer surface of the cell membrane protects the cell membrane by detoxifying free radicals. Oxygen (O₂), hydrogen peroxide (H₂O₂), and hydroxy (OH) formed during mitochondrial respiration reduce the release of free radicals.

Provides protection against the oxidation of DNA, molecule is not easily oxidized due to its lack of peroxidation activity. In addition, it does not enter the redox cycle and radical forming reactions [20]. Unlike other antioxidants, it has no toxic effect when used in excessive amounts. The various classic antioxidants (vitamin E, vitamin C, beta-carotene, etc.) become pro-oxidants once they show their action. These substances are less harmful than the oxidants they remove. However, melatonin is an oxidant, and its by-products also have antioxidant effects. Its property is very valuable for an antioxidant agent and is described as a 'terminal antioxidant. In conclusion; Melatonin has been found to stimulate antioxidant enzymes, prevent lipid peroxidation, and protect brain tissue from free radicals produced by oxygen intake [21-22].

Melatonin and the digestive system. Melatonin is synthesized in high concentrations in enter chromaffin cells of the gastrointestinal tract. Not only the epiphyseal enzyme of melatonin synthesis hydroxyl-O-methyltransferase (HIOMT) and its precursor serotonin are found in enter chromaffin cells of the intestinal mucosa. Increases bicarbonate (HCO₃) secretion via melatonin receptors (MT₂) in duodenal epithelial cells and enhances the protective factor.

Melatonin acts in the digestive system in addition to its receptor effect as well as in a non-receptor way. By removing free radicals, GI has an inhibitory effect on ulcer formation [24].

Effects of melatonin on digestive diseases. Melatonin is known to be synthesized in significant amounts in the gastric mucosa. Because of its lyophilic properties, the results of immunohistochemically studies support the hypothesis that it can penetrate deeper into the mucosal layer, even into the intermuscular plexus, and into the muscular layers [28]. The gastric mucosa is constantly exposed to factors such as hydrochloric acid and pepsin, which can damage the tissue. There are many mechanisms that protect the structure and function of the stomach against these factors, such as the mucosal bicarbonate barrier and prostaglandins. An imbalance between the harmful agents and the protective mechanisms causes damage to the stomach.

Gastric damage or ulceration has been induced in experimental animal studies by various means, such as ethanol administration, injection of indomethacin, exercise restriction, and cold stress [25]. Studies carried out in the early 2000s showed that stress-induced gastric ulcer formation in an ischemia-reperfusion model was suppressed by the administration of melatonin and L-tryptophan. These studies have suggested that the therapeutic and prophylactic effects of melatonin on gastric mucosal damage are based on stimulation of the cyclooxygenase pathway, an increase in prostaglandin

synthesis, and induction of nitric oxide (NO) synthesis. It has also been shown to accelerate blood flow in the mechanism of action.

These results are consistent with studies showing an increase in inducible nitric oxide synthase (iNOS) mRNA and its expression at the edges of healing ulcers [26,27].

The role of the antioxidant effect of melatonin through the removal of reactive oxygen species in this protective and therapeutic effect cannot be excluded. In fact, animal studies in which cyclooxygenase enzymes and prostaglandin synthesis are inhibited by indomethacin have shown that melatonin has therapeutic and protective effects due to its antioxidant activity.

In addition to all these effects, there is evidence that the increased concentrations of gastrin and cholecystokinin observed in patients treated with melatonin and L-tryptophan also promote ulcer healing [29-31]. In light of these studies, the ability of melatonin and L-tryptophan to treat gastric mucosal damage can be summarized. Other important pathologies such as gastroesophageal reflux disease (GERD), chronic esophagitis, Barrett's esophagus, and esophageal cancer are common in the community. The incidence of Barrett's esophagus and esophageal strictures is increasing. For this reason, the effect of melatonin in patients with gastroesophageal reflux disease has been specifically studied. One of these, melatonin and L-tryptophan, has been clinically tested for reducing the symptoms of gastroesophageal reflux disease and achieving complete remission [32]. In an animal model of gastroesophageal reflux disease, prior administration of melatonin prevented esophageal damage. In the same study, it was shown that increased mucosal blood flow, increased prostaglandin synthesis (PGE₂), and decreased TNF-alpha levels were associated with a therapeutic effect of melatonin [33]. Reduced plasma melatonin levels in patients with digestive diseases, such as gastroesophageal reflux disease and duodenal ulcer disease, have shown that deficiency of this hormone influences the rapid damage of the upper gastrointestinal mucosa. Reduced melatonin synthesis in the elderly patient group and a higher risk of complications due to reflux esophagitis in elderly patients support the possibility of a possible harmful effect of melatonin deficiency on upper gastrointestinal mucosa [33,34].

Inflammatory bowel diseases, more commonly Crohn's disease and ulcerative colitis, are chronic inflammatory pathologies with remissions with relapses and temporary improvements. Triggers are also the occurrence of chronic inflammation, the presence of some genetic predisposition, and environmental factors that compromise the integrity of the intestinal epithelial barrier. There are studies that show that circadian rhythm influences the composition of the intestinal microbiota. There may be a hypothetical link between inflammatory bowel disease and melatonin secretion. A clinical trial reported that melatonin significantly improved outcomes in Crohn's disease and ulcerative colitis when prescribed in addition to basic treatment for inflammatory bowel disease [25].

By reducing the expression of melatonin-induced nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) in the colon, acetyl acid enemas with 2,4,6-trinitrobenzene sulphonic acid (TNBS) have been shown to have a

protective effect on the intestinal mucosa against models of induced colitis [26].

The role of melatonin in clinical use. Today, melatonin and its agonists have found use in clinical practice. Melatonin medications are available with or without a prescription. Melatonin is used for sleep disorders, a syndrome of circadian rhythm disturbance due to short journeys called 'jetlag' [27]. On the other hand, drugs such as ramelteon, which are melatonin agonists, are indicated for insomnia, and agomelatine is indicated for depression associated with sleep disorders [28,29]. Current applications of melatonin are depression, including biological rhythm and sleep regulation.

Aim. To monitor the therapeutic effects of melatonin in relation to ethanol concentration in an ethanol-induced experimental gastric ulcer model.

Materials and methods

Study: Experimental (preclinical). Preclinical model details.

Subjects: Wistar Albino rats (300-380 g).

Induction of Ulcers: Gastric ulcers induced using ethanol at varying concentrations (100%, 75%, 60%). This concentrations of ethanol through a special gastric probe and waited for 1 hour to develop an ulcer model.

Groups: group 1: Ethanol-only (control). n=6;

group 2 (pre - melatonin): Ethanol + melatonin (15 mg/kg) administered intraperitoneally. Melatonin (15 mg/kg) was administered intraperitoneally before 30 min ethanol consumption. n=6;

group 3 (pre - melatonin): Control group. Melatonin (15 mg/kg) was administered intraperitoneally before 30min euthanasia n=6;

Melatonin dosage: Administered intraperitoneally at 15 mg/kg.

Outcome Measures: Macroscopic analysis of gastric lesions (ulcer area).

Etics approval. Approval for the research was received from Gazi University Animal Experiments Local Ethics Committee (Number: 04.10.2019-E.123689).

The study was conducted in group 6, with models of gastric ulcer induced by ethanol in different concentrations (100-75-60%) and pre-melatonin (15 mg/kg) administered intraperitoneally in the ulcer group and control group. This preclinical study was conducted in the research laboratory of the Department of Clinical Pharmacology, Faculty of Medicine, Gazi University, Turkey. Ethanol-induced gastric ulcer model Rats weighing 300-380 g, matured under standard conditions (Wistar Albino breed), were injected with different concentrations of ethanol through a special gastric probe and waited for 1 hour to develop an ulcer model. Subsequently, euthanasia was carried out under anesthesia, the stomach was extracted and macroscopic analysis was performed. The wound area was measured using a millimeter ruler. Statistically, variables were summarized using frequencies and percentages. $P < 0.05$ was considered statistically significant.

The results

In the rat stomach, ethanol increased the wound area in a concentration-dependent macroscopic analysis. In addition to this increase in the wound area, melatonin administered as a prophylactic subcutaneous injection reduced the total wound area. In the ethanol-induced group (100%), the wound area was 15.01 ± 1.92 mm, with melatonin (15 mg/kg) administered intraperitoneally, and the wound area was 14.01 ± 1.74 mm after 30 minutes in the 100% ethanol-induced group, 13.24 ± 1.71 mm in the 75% ethanol group, 11.87 ± 1.45 mm in the melatonin + 75% ethanol induced group, the wound area was 11.52 ± 1.15 mm in the 60% ethanol induced group and 6.12 ± 0.54 mm in the melatonin + 60% ethanol induced group [Table 1; Figure 1,2].

Table 1.

Effects of melatonin in a model of gastric ulcer induced by ethanol at different concentrations.

Concentration	n	Macroscopic wound area	Concentration	n	Macroscopic wound area
Ethanol 100%	6	15.01 ± 1.92	Melatonin + Ethanol %	6	14.01 ± 1.74
Ethanol 75%	6	13.24 ± 1.71	Melatonin + Ethanol 75%	6	11.87 ± 1.45
Ethanol 60%	6	11.52 ± 1.15	Melatonin + Ethanol 60%	6	6.12 ± 0.54



Figure 1. Model of an ethanol-induced gastric ulcer.



Figure 2. Melatonin + ethanol gastric ulcer model.

Rats in the control group were put to sleep by anesthesia after 24 hours of fasting without any manipulation. In the control group, no signs of gastric mucosal lesions were observed when the gastric mucosa was analyzed. 1 h after administration of ethanol through the gastric tube, the mucosa of the group which developed gastric ulcer showed erosions, ulcers, hemorrhages, petechial with black spots, puffiness, swelling of the mucosa, and an area of lesions with redness. 1 hour later. 30 min after intraperitoneal injection of melatonin (15 mg/kg) in rats administered ethanol via a gastric tube, the mucosal layer showed erosions, reduced ulcerated area, pulmonary petechial, edema, the reddening area of the mucosal layer was statistically reduced compared with the wound model group, and no bleeding was observed.

Discussion

A single blind randomized study was performed in which 176 patients underwent treatment using the supplement cited and 175 received treatment of 20 mg omeprazole. All patients with melatonin (100%) reported a complete regression of symptoms after 40 days of treatment. On the other hand, 115 subjects (65.7%) of the omeprazole reported regression of symptoms in the same period. There was statistically significant difference between the groups ($P < 0.05$). This formulation promotes regression of GERD symptoms with no significant side effects [22].

In a gastric ulcer model, it has been shown that prior administration of melatonin through the gastric tube reduces the area of ulcer formation. In many experimental and clinical studies, melatonin pre-injection has been shown to reduce pathological and inflammatory processes in various models of the digestive system. When the ethanol concentration is high, the wound area is enlarged and bleeding is more severe, while in the groups receiving lower concentrations, the wound area is relatively reduced compared to the 100% concentration. The therapeutic properties of melatonin depend on the concentration of ethanol. In other words, melatonin exhibited a therapeutic effect to varying degrees, depending on the strength of the aggressive factor. Particularly in the group of wounds induced by 60% ethanol, the therapeutic protective effect of melatonin was statistically significant. Melatonin has been shown in many studies to reduce ulcer index and lesion area in a model of gastric ulcer induced by other factors such as ethanol, stress, indomethacin, and non-steroidal anti-inflammatory drugs. It has been suggested that the mechanism of action lies in the stimulation of nitric oxide and prostaglandin synthesis. In addition, melatonin has been shown to exert its effects by increasing mRNA levels and the expression of inhibitory nitric oxide synthase (iNOS) in the tissue surrounding the wound. In a model of gastroesophageal reflux disease, its beneficial and protective effects on the esophageal mucosa were attributed to its antioxidant and anti-inflammatory effects. It has been shown to reduce the secretion of pro cytokines involved in inflammation, such as IL-1 β and TNF- α [29-31].

Application of melatonin abrogates mesothelial cell pyroptosis through a MT1R-mediated action, and successfully reduces peritoneal fibrosis and angiogenesis in an animal model while preserving dialysis efficacy. Mechanistically, melatonin treatment maintains mitochondrial integrity in mesothelial cells, meanwhile

activating mTOR signaling through an increase in the glycolysis product dihydroxyacetone phosphate [32].

In a clinical trial, the use of melatonin alone and in combination with omeprazole in 36 patients diagnosed with gastroesophageal reflux disease showed very significant results in reducing complaints [33].

Conclusion

Melatonin is known to be an important chronobiological molecule that exerts its effect on any system in the body. The role and pharmacological effect of melatonin in the gastrointestinal system is still one of the important issues, and research continues in various directions. In this study, melatonin showed a positive effect in the prevention and treatment of ethanol-induced ulcers at different concentrations. A significant change was observed at higher levels of melatonin inhibition of ethanol exposure at lower concentrations. We hope that in clinical practice, taking into account its therapeutic effect on the mucous membrane of the digestive system through its antioxidant effect against oxidative stress, and taking into account the preventive and therapeutic properties of the digestive system, the indicator of wide-spectrum use will increase. For this reason, it is necessary to conduct clinical studies in different directions for the use of its antioxidant effect.

Etics approval.

Approval for the research was received from Gazi University Animal Experiments Local Ethics Committee (Number: 04.10.2019-E.123689).

Authors' contributions (распишите Фамилию и инициал)

Concept: Skenderova A.A., Nuskabayeva G.O., Tatykayeva U.B., Sarsenbayeva G.Zh., Kemelbekov K.S., Seidakhmetova A.A.,

Final approval: All listed authors.

This study was presented at the 26th National and 1st International Pharmacology Congress [34].

Conflicts of interest: All authors declare no conflict of interest.

Funding. This study was supported by Gazi University BAP Unit with project number 01.2019-59.

References:

1. Bubenik George A. Gastrointestinal melatonin: localization, function, and clinical relevance. Digestive diseases and sciences. 2002. №47. P. 2336-2348. DOI:10.1023/a:1020107915919
2. Boutin J.A., Kennaway D.J., Jockers R. Melatonin: Facts, Extrapolations and Clinical Trials. Biomolecules, 2023. 13(6), 943. <https://doi.org/10.3390/biom13060943>
3. Brzozowski Tomasz et al. The role of melatonin and L-tryptophan in prevention of acute gastric lesions induced by stress, ethanol, ischemia, and aspirin. Journal of pineal research. 1997. №23(2). P.79-89. DOI:10.1111/j.1600-079x.1997.tb00339.x
4. Carrillo-Vico Antonio et al. Evidence of melatonin synthesis by human lymphocytes and its physiological significance: possible role as intracrine, autocrine, and/or paracrine substance. The FASEB Journal. 2004. №18.3. P. 537-539. DOI:10.1096/fj.03-0694fje
5. Claustrat Bruno, Brun Jocelyne, Chazot Guy. The basic physiology and pathophysiology of melatonin. Sleep medicine reviews, 2005. №9(1). P. 11-24. DOI:10.1016/j.smrv.2004.08.001

6. Conway S, Canning SJ, Howell HE, Mowat ES, Barrett P, Drew JE, et al. Characterisation of human melatonin mt(1) and MT(2) receptors by CRE-luciferase reporter assay. *Eur J Pharmacol* 2000. 390:15-24.
7. Cruz-Sanabria F., Carmassi C., Bruno S., Bazzani A., Carli M., Scarselli M., Faraguna U. Melatonin as a Chronobiotic with Sleep-promoting Properties. *Current neuropharmacology*, 2023. 21(4), 951-987. <https://doi.org/10.2174/1570159X20666220217152617>
8. Choudhary P., Roy T., Chatterjee A., Mishra V.K., Pant S., Swarnakar S. Melatonin rescues swim stress induced gastric ulceration by inhibiting matrix metalloproteinase-3 via down-regulation of inflammatory signaling cascade. *Life sciences*, 2022. 297, 120426. <https://doi.org/10.1016/j.lfs.2022.120426>
9. Chang C.L., Chen C.H., Chiang J.Y., Sun C.K., Chen Y.L., et al. Synergistic effect of combined melatonin and adipose-derived mesenchymal stem cell (ADMSC)-derived exosomes on amelioration of dextran sulfate sodium (DSS)-induced acute colitis. *American journal of translational research*, 2019. 11(5), 2706-2724. PMID: 31217848
10. Faghih Dinevari M., Jafarzadeh F., Jabbaripour Sarmadian A., Abbasian S., Nikniaz Z., Riaz A. The effect of melatonin on irritable bowel syndrome patients with and without sleep disorders: a randomized double-blinded placebo -controlled trial study. *BMC gastroenterology*, 2023. 23(1), 135. <https://doi.org/10.1186/s12876-023-02760-0>
11. Grigg-Damberge, Madeleine M., Ianakieva Dessislava. Poor quality control of over-the-counter melatonin: what they say is often not what you get. *Journal of Clinical Sleep Medicine*. 2017. №13(2). P. 163-165. DOI: 10.5664/jcsm.6434
12. Hardeland R. Redox Biology of Melatonin: Discriminating Between Circadian and Noncircadian Functions. *Antioxidants & redox signaling*, 2022. 37(10-12), 704-725. <https://doi.org/10.1089/ars.2021.0275>
13. Jaworek Jolanta, Brzozowski Tomasz, Konturek Stanislaw Jan. Melatonin as an organoprotector in the stomach and the pancreas. *Journal of pineal research*. 2005. №38(2). P.73-83. DOI: 10.1111/j.1600-079X.2004.00179.x
14. Kato K. et al. Circadian rhythm of melatonin and prostaglandin in modulation of stress-induced gastric mucosal lesions in rats. *Alimentary pharmacology & therapeutics*. 2002. №16. P.29-34. DOI: 10.1046/j.1365-2036.16.s2.11.x
15. Konturek Peter C. et al. Esophagoprotection mediated by exogenous and endogenous melatonin in an experimental model of reflux esophagitis. *Journal of pineal research*. 2013. № 55 (1). P. 46-57. DOI:10.1111/jpi.12048
16. Konturek S.J. et al. Localization and biological activities of melatonin. *J. Physiol. Pharmacol*. 2007. №58(3) P. 381-405. PMID:17928638
17. Konturek S.J., et al. Protective influence of melatonin against acute esophageal lesions involves prostaglandins, nitric oxide and sensory nerves. *Journal of Physiology and Pharmacology*. 2007. №58(2). P. 361. PMID:17622703
18. Mayo J.C., Sainz R.M. Melatonin from an Antioxidant to a Classic Hormone or a Tissue Factor: Experimental and Clinical Aspects 2019. *International journal of molecular sciences*, 2020. 21(10), 3645. <https://doi.org/10.3390/ijms21103645>
19. Macchi M.M., Bruce J.N. Human pineal physiology and functional significance of melatonin. *Front Neuroendocrinol*. 2004. № 25. P. 177-195. <https://doi.org/10.1016/j.yfrne.2004.08.001>
20. Majka, Jolanta, et al. Melatonin in prevention of the sequence from reflux esophagitis to Barrett's esophagus and esophageal adenocarcinoma: experimental and clinical perspectives. *International journal of molecular sciences* - 2018. №19(7). P. 2033. DOI: 10.3390/ijms19072033
21. Michurina S.V., Kolesnikov S.I., Ishchenko I.Y., Arkhipov S.A. Light-Induced Functional Pinealectomy: Expression of MT2 Receptors in Liver Cells of C57BL/6 Mice after Melatonin Treatment. *Bulletin of experimental biology and medicine*, 2022. 173(4), 569-574. <https://doi.org/10.1007/s10517-022-05590-4>
22. Mehrzadi S., Sheibani M., Koosha F., Alinaghian N., Pourhanifeh M.H., Tabaeian S.A.P., Reiter R.J., Hosseinzadeh A. Protective and therapeutic potential of melatonin against intestinal diseases: updated review of current data based on molecular mechanisms. *Expert review of gastroenterology & hepatology*, 2023. 17(10), 1011-1029. <https://doi.org/10.1080/17474124.2023.2267439>
23. Özçelik Fatih et al. Melatonin: Genel özellikleri ve psikiyatrik bozukluklardaki rolü. *Psikiyatride Güncel Yaklaşımlar*. 2013. №5(2). P. 179-203. <https://doi.org/10.5455/cap.20130512>
24. Pereira Ricardo de Souza. Regression of gastroesophageal reflux disease symptoms using dietary supplementation with melatonin, vitamins and aminoacids: comparison with omeprazole. *Journal of Pineal Research*. 2006. №41(3). P. 195-200. DOI:10.1111/j.1600-079X.2006.00359.x
25. Potes Y., Caballero B. 2023. Special Issue on "Pleiotropic Benefits of Melatonin: From Basic Mechanisms to Disease". *International journal of molecular sciences*, 24(6), 5223. <https://doi.org/10.3390/ijms24065223>
26. Peña-Mercado E., Garcia-Lorenzana M., Huerta-Yepez S., Cruz-Ledesma A., Beltran-Vargas N.E. Effect of melatonin on electrical impedance and biomarkers of damage in a gastric ischemia/reperfusion model. *PloS one*, 2022. 17(8), e0273099. <https://doi.org/10.1371/journal.pone.0273099>
27. Ratzburg Courtney. Melatonin-the myths and facts. *Vanderbilt University*. 2007. №12(02). P.376-384
28. Ruan H., Li X., Zhou L., Zheng Z., Hua R., Wang X., Wang Y., Fan Y., et al. Melatonin decreases GSDME mediated mesothelial cell pyroptosis and prevents peritoneal fibrosis and ultrafiltration failure. *Science China. Life sciences*, 2024. 67(2), 360-378. <https://doi.org/10.1007/s11427-022-2365-1>
29. Sahoo D.K., Heilmann R.M., Paital B., Patel A., Yadav V.K., Wong D., Jergens A.E. Oxidative stress, hormones, and effects of natural antioxidants on intestinal inflammation in inflammatory bowel disease. *Frontiers in endocrinology*, 2023. 14, 1217165. <https://doi.org/10.3389/fendo.2023.1217165>
30. Skenderova A., Sebile Koç D., Güvel M.C., Öztürk Fincan G., Uluoğlu C. The role of melatonin and nitric oxide in the neurogenic contraction responses of rat fundus strips

to electrical field stimulation in the peptic ulcer model. 26th National and 1st International Pharmacology Congress; Turkish Pharmacology Association, Turkey, 2021. P.1264-1271

31. Tassan Mazzocco M., Pisanu C., Russo L., Acconcia, C., Cambiaghi M., De Girolamo S., Squassina A., Cherchi L. 2023. Melatonin MT₁ receptors as a target for the psychopharmacology of bipolar disorder: A translational study. *Pharmacological research*, 198, 106993. <https://doi.org/10.1016/j.phrs.2023.106993>

32. Tordjman Sylvie et al. Melatonin: pharmacology, functions and therapeutic benefits. *Current*

neuropharmacology. 2017. № 15(3). P. 434-443. DOI:10.2174/1570159X14666161228122115

33. Yazıcı Cevat, Kader Köse. Melatonin: karanlığın antioksidan gücü. *Sağlık Bilimleri Dergisi*. 2004. №13(2). P. 56-66.

34. Zagajewski J., Wojcik-Grzybek D., Brzozowski B., Majka J., et al. Simultaneous detection of melatonin and six metabolites of L-tryptophan pathway in rat gastric mucosa. *Journal of physiology and pharmacology: an official journal of the Polish Physiological Society*, 2021. 72(6), 10.26402/jpp.2021.6.15. <https://doi.org/10.26402/jpp.2021.6.15>

Information about authors:

Akbota A. Skenderova – PhD, Senior Lecturer, Department of Internal Medicine, num: 87784884285, e-mail: akbota.skenderova@ayu.edu.kz; <https://orcid.org/0000-0002-5392-9433>, «Akhmet Yassawi International Kazakh-Turkish University», Turkistan, Republic of Kazakhstan.

Gulnaz O. Nuskabayeva Candidate of Medical Sciences, Associate Professor, Department of Specific Clinical Disciplines num: 87052853131, e-mail: nuskabayeva.gulnaz@ayu.edu.kz; <https://orcid.org/0000-0002-2906-1427>, «Akhmet Yassawi International Kazakh-Turkish University», Turkistan, Republic of Kazakhstan.

Ugilzhan B. Tatykayeva, Candidate of Medical Sciences, Senior Lecturer, Department of Fundamental Medical Disciplines, num: 87026141701, e-mail: ugilzhan.tatykayeva@ayu.edu.kz; <https://orcid.org/0000-0002-7519-479X>, «Akhmet Yassawi International Kazakh-Turkish University», Turkistan, Republic of Kazakhstan.

Gulzat Zh. Sarsenbayeva – Candidate of Medical Sciences, Associate Professor, Head of the Department “Social health insurance and public health”, num: 87017496288, e-mail: gulzat.zh@mail.ru; <https://orcid.org/0000-0002-1518-6528>, JSC South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan;

Kanat-zhan S. Kemelbekov – PhD., Head of the Department “Pediatrics-1, num: 87085334300, e-mail: kanat-270184@mail.ru; <https://orcid.org/0009-0008-1619-6278>, JSC South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan;

Aizat A. Seidakhmetova - Candidate of Medical Sciences, Head of the Department of "Emergency Medicine and Nursing", num: 87017671448, e-mail: aizat67@mail.ru; <https://orcid.org/0000-0002-5794-3814>, JSC South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan;

Corresponding author:

Akbota Skenderova, PhD, Department of General Surgery, Akhmet Yassawi International Kazakh-Turkish University, Turkistan city, Republic of Kazakhstan.

Postal address: 161200, Republic of Kazakhstan, Turkestan, Bekzat Sattarkhanov Avenue, 29/3

E-mail: akbota.skenderova@ayu.edu.kz

Phone: +7 778 488 42 85.

